

# A new species of the cicada genus *Cicadatra* Kolenati, 1857 (Hemiptera, Cicadidae) from Pakistan with a key to the known species of Pakistani *Cicadatra*

Zubair Ahmed<sup>1,†</sup>, Allen F. Sanborn<sup>2,‡</sup>, Muhammad Atique Akhter<sup>3,§</sup>

- 1** Department of Zoology, Federal Urdu University of Arts, Sciences & Technology, Karachi, Pakistan  
**2** Department of Biology, Barry University, 11300 NE Second Avenue. Miami Shores, FL 33161-6695, USA  
**3** Department of Zoology, University of Karachi, Karachi, Pakistan

† [urn:lsid:zoobank.org:author:77ABB331-ACE3-4A2E-959C-A7E7CCF77686](https://doi.org/urn:lsid:zoobank.org:author:77ABB331-ACE3-4A2E-959C-A7E7CCF77686)

‡ [urn:lsid:zoobank.org:author:9544FF3F-9B1F-457A-90A2-8A1C0C1F22A0](https://doi.org/urn:lsid:zoobank.org:author:9544FF3F-9B1F-457A-90A2-8A1C0C1F22A0)

§ [urn:lsid:zoobank.org:author:B00AE5E6-D0ED-4062-ABBD-A74BD44FAFD2](https://doi.org/urn:lsid:zoobank.org:author:B00AE5E6-D0ED-4062-ABBD-A74BD44FAFD2)

Corresponding author: Zubair Ahmed ([zubair\\_ahmed\\_74@yahoo.com](mailto:zubair_ahmed_74@yahoo.com))

---

Academic editor: M. Wilson | Received 31 October 2011 | Accepted 27 February 2012 | Published 9 March 2012

[urn:lsid:zoobank.org:pub:DB67F6F7-2BC1-4D26-8193-00C30487FBE9](https://doi.org/urn:lsid:zoobank.org:pub:DB67F6F7-2BC1-4D26-8193-00C30487FBE9)

---

**Citation:** Ahmed Z, Sanborn AF, Akhter MA (2012) A new species of the cicada genus *Cicadatra* Kolenati, 1857 (Hemiptera, Cicadidae) from Pakistan with a key to the known species of Pakistani *Cicadatra*. ZooKeys 174: 41–48. doi: 10.3897/zookeys.174.2299

---

## Abstract

A new species of cicada, *Cicadatra zianatica* **sp. n.**, is described from Pakistan. Male genitalia, timbal and opercula are described and illustrated as important diagnostic characters. Biological notes are also provided. A key to the known *Cicadatra* of Pakistan is provided.

## Keywords

*Cicadatra* species, taxonomy, Cicadidae, morphology

## Introduction

Species of the genus *Cicadatra* Kolenati, 1857 exhibit variability in several morphological characters. As a result, there is still much to learn about the distribution and composition of *Cicadatra* species across the range of the genus. Studies of *Cicadatra* species in the Middle East and Asia continue to illustrate the lack of knowledge about

the species of this widespread genus. However, there have been some more recent studies which provide detailed analyses of the morphological characters of generally new species from this region (Dlabola 1960, 1970, 1979, 1981, Dlabola and Heller 1962, Linnavouri 1962, Boulard 1977, Melichar 1896, Mirzayans 1995, Mozaffarian and Sanborn 2010, Mozaffarian et al. 2010, Ahmed and Sanborn 2010, Ahmed et al. 2010) including the first checklists of the genus for Iran (Mozaffarian and Sanborn 2010) and Pakistan (Ahmed and Sanborn 2010).

The first checklist of the Pakistani cicada fauna was recently produced by Ahmed and Sanborn (2010). They determined that 29 known species inhabited Pakistan at the time with the first records of seven species and the descriptions of four new species included in the total (Ahmed and Sanborn 2010). They listed seven species of *Cicadatra* for Pakistan, four of which represented new records for the country. Later that same year, Ahmed et al. (2010) described a new species of *Cicadatra* from Pakistan along with sound analysis and DNA sequencing of the species. The current work represents another new species that was collected during recent fieldwork. A more complete knowledge of the Pakistani cicada fauna will only be obtained with continued field research and studies of existing collections.

The present species is described from Pakistan as new to science, and is known only from the Ziarat District, Balochistan Province. Notes on the biology of this new species are also provided along with a key to differentiate the species from the known Pakistani *Cicadatra*.

## Materials and methods

Specimens were captured during June 2010 and June 2011 in Balochistan Province, Pakistan. Terminology follows Moulds (2005). Measurements were made with Vernier calipers or a Wild Heerbrugg 12034 binocular microscope. Specimens are deposited in the collections of the Natural History Museum, University of Karachi, Pakistan (NHMK) and Zubair Ahmed Collection, Pakistan (ZACP).

## Results and discussion

### *Cicadatra ziaratica* Ahmed, Sanborn & Akhter, sp. n.

urn:lsid:zoobank.org:act:04C75272-8CDE-4677-A6EE-798512C83B53

[http://species-id.net/wiki/Cicadatra\\_ziaratica](http://species-id.net/wiki/Cicadatra_ziaratica)

Figs 1–8

**Type locality.** Pakistan, Balochistan Province, Khotal Chehri, District Ziarat.


**Type specimens.** Holotype male, pinned. Original label: “Pakistan, Balochistan Province, Khotal Chehri, District Ziarat, 7.vi.2010, Collector, Zubair Ahmed”, “HOLOTYPE / *Cicadatra ziaratica* / Ahmed, Sanborn & Akhter” [handwritten label]



**Figure 1.** Holotype Male, *Cicadatra ziaratica* sp. n.

(NHMK); one male paratype, “Pakistan, Balochistan Province, Khotal Chehri, District Ziarat, 7.vi.2010, Collector, Zubair Ahmed”, “PARATYPE / *Cicadatra ziaratica* / Ahmed, Sanborn & Akhter” [handwritten label]; three male paratypes, “Pakistan, Balochistan Province, Khotal Chehri, District Ziarat, N, 3.vi.2011, Collector, Zubair Ahmed “PARATYPE / *Cicadatra ziaratica* / Ahmed, Sanborn & Akhter” [handwritten label] (ZACP).

**Diagnosis.** The new species appears to be most allied morphologically to *C. lorestanica* Mozaffarian and Sanborn 2010 from Iran and *C. karachiensis* Ahmed et al. 2010 from Pakistan. The new species can be distinguished by the upper lobe of the pygofer being ill-defined in *C. ziaratica* whereas it is a finger-like extension in *C. lorestanica*. The aedeagus of *C. lorestanica* has a curved, bifold, sclerized, hook-like process and two lateral spiny appendages while the aedeagus of *C. ziaratica* has a long, subapical spine, a dorsal spine along sclerized teeth-like process, a lateral spine and a ventral semicircular toothed process. In *C. karachiensis* the upper lobe of the pygofer is rounded, the aedeagus has a long upturned flap with 11 aedeagal spines and the hind wing has five apical cells instead of the six found in *C. ziaratica*. The species are similar in possessing a mesonotum with two lines but the degree of curvature is slightly variable. Fore wings with radial and radiomedial crossveins at bases of the 2<sup>nd</sup> and 3<sup>rd</sup>



**Figures 2–8.** *Cicadatra ziaratica* sp. n., 2 Male pygofer lateral view 3–4 Aedeagus 5 Claspers 6 Timbal cover 7 Timbal 8 Operculum. Scale lines = 0.6 mm. **Asf** Aedeagus with serrated flap **Als** Aedeagus long spine **Ls** Lateral spine **Rss** Row of middle spines

apical cells infuscated in *C. ziaratica* but lacking infuscation in *C. lorestanica* and *C. karachiensis*. The timbal cover of *C. karachiensis* is reduced and ventral to the majority of the timbal while the timbal cover in *C. ziaratica* and *C. lorestanica* covers more than

half the timbal and is centrally located over the timbal. Finally, the timbal has 9 ribs in *C. ziaratica* and *C. karachiensis* but 11 ribs in *C. lorestanica*. The remaining species known to inhabit Pakistan can be distinguished using the key. There are insufficient data to perform a molecular phylogenetic analysis of the Pakistani *Cicadatra* species as genes from a limited number of species have been sequenced (Ahmed et al. 2010).

**Description. Male.** General color of body black with olive to ochraceous markings and white pile.

Head black with white pile particularly on posterior edge, head including eyes as broad as mesonotum; eyes brown, varying from light to dark in different specimens; ocelli orangish, piceous in some specimens; postclypeus black with a central sulcus and obvious transverse grooves, dense pile lateral of grooves, gena and lorum black with dense white pile; rostrum light ochraceous at base, darker towards apex, strongly passing intermediate coxae; labrum with sparse white pile laterally and on apex; antennae dark brown, apical segment faint yellow, vertex black, supra-antennal plate reaching eyes, black, light band at medially in two paratypes.

Pronotum black, brown in some paratypes, with median black biconcave mark containing a light olive green median fascia, an olive green patch crossing ambient fissure posterolateral to each side of median fascia, black mark continues around disc in ambient fissure and across lateral pronotal collar to the lateral angle; paramedian and lateral fissures variably marked with dark brown to black, pronotal collar black anteriorly and olive green across posterior half of lateral angles and posterior margin, ochraceous in some paratypes, white dense pile present on ambient, paramedian and lateral fissure and scattered pile on disc, pile reduced in some paratypes; mesonotum black or dark brown, with ochraceous J-shaped mark along parapsidal suture, mark triangularly shaped at base in some paratypes; cruciform elevation olive green (brown in some paratypes) medially, darkening to black in anterior arms; metanotum olive green (brown in some paratypes); thoracic sternites black with dense white pile, ochraceous in different specimens; some specimens with dark marking on basisternum 2, epimeron 2, katapisternum 2, and episternum 3.

Fore coxae light olive to ochraceous with black linear marking, middle coxae light ochraceous with broad dark anterolateral surface, hind coxae light ochraceous with darker laterally; fore and middle trochanter olive to ochraceous with a dark brown area at middle with white pile; fore femorae dark brown with white pile and light areas on ventral apex with strongly angled primary spine, erect secondary spine and a small angled apical spine; middle femora dark brown with a yellow area at ventral and apex with dense white pile, hind femora dark brown with yellow area on base and apex; fore tibiae dark brown lighter at apex, middle tibiae dark brown with white pile and yellow at lateral, hind tibiae yellow, half dark brown with five brown tibial spurs and sparse white pile; tibial spurs and combs brown, darker towards their apices; tarsi black; pre-tarsal claws dark brown.

Fore wings hyaline with faint yellow and brown venation, radial (r) and radiomedial (r-m) crossveins at bases of apical cells 2 and 3 darkly infuscated, infuscation on r-m absent or reduced in some paratypes, basal call twice as long as wide; fore wings

with 8 apical cells, basal membrane light reddish; hind wings with faint yellow venation, light grey infuscation around anal veins 2 and 3 (2A and 3A), hind wings with 6 apical cells.

Male opercula light brown with black spot on lateral base and rather dense white pile, rounded, and slightly overlapped, not meeting medially in paratypes, meracanthus triangular, light ochraceous with black spot at base.

Abdominal tergites black with white pile more or less located near the anterior edge of each tergum, tergites 2–7 with a light area on posterior except median part, timbal cavity exposed; timbal cover incomplete covering about half the timbal, black or dark brown with white pile, timbal with 9 ribs; abdominal sternites brown with dense white pile, epipleurites dark brown with dense white pile.

Male pygofer dark brown with scattered pile, dorsal beak pointed, upper lobe of pygofer rounded, basal lobe of pygofer appears as a bud like projection beneath the upper lobe; uncus very short; claspers tapering to a point, curved slightly laterad, close to each other at base; aedeagus with theca curved, a lateral scleritized, serrate appendages, a long, subapical spine, a ventral scleritized, rounded serrate process, a lateral and a median long spine.

**Female.** Unknown.

**Etymology.** The species is named for the district of Balochistan from which the type series was collected.

**Measurements (mm).** N=5 males, mean (range). Length of body: 16.9 (16.0–18.0); length of fore wing: 20.2 (19.0–22.0); width of fore wing: 6.7 (6.1–7.0); width of head including eyes: 4.6 (4.0–5.0); width of pronotum including paratota: 5.7 (5.5–6.0); width of mesonotum: 5.1 (4.8–5.5).

**Biological notes.** All specimens were collected during 2010 and 2011 in the vicinity of Ziarat between 3 June–7 June. The cicadas emerged among wild grasses based on the location of the emergence holes. Adult males called from these same grasses as well as from shrubs including *Peganum harmala* L.

### Key to the males of *Cicadatra* of Pakistan

- 1 Body length >24 mm, pronotal collar almost or entirely black ..... 2
- Body length <24 mm, pronotal collar heavily marked with ochraceous, olivaceous or tawny ..... 4
- 2 Pronotal disk black ..... *C. acberi* (Distant, 1888)
- Pronotal disk ochraceous ..... 3
- 3 Supra-antennal plate, cruciform elevation, and costal margin tawny .....  
..... *C. persica* Kirkaldy, 1909
- Supra-antennal plate and cruciform elevation black, costal margin castaneous ..... *C. gingat* China, 1926
- 4 Head castaneous ..... *C. raja* (Distant, 1906)
- Head black ..... 5

5 Radial and radiomedial crossveins not infuscated..... 6  
 – Radial and/or radiomedial crossveins infuscated..... 7  
 6 Pronotum castaneous marked with black, cruciform elevation black, male opercula overlapping medially, small marginal spot on hind wing.....  
 ..... ***C. sankara* (Distant, 1904)**  
 – Pronotum dark ochraceous marked with black, cruciform elevation marked with ochraceous, male opercula almost meeting medially, hind wing hyaline ..... ***C. karachiensis* Ahmed, Sanborn and Hill, 2010**  
 7 Postclypeus black ..... ***C. ziaratica* sp. n.**  
 – Postclypeus tawny marked with castaneous or black ..... **8**  
 8 Postclypeus with transverse grooves black ..... ***C. walkeri* Metcalf, 1963**  
 – Postclypeus with medial castaneous or black stripe.....  
 ..... ***C. xanthes* (Walker, 1850)**

**Acknowledgements**

ZA thanks Tim McNary, Colorado USA for providing pertinent literature. Collection of specimens was partially supported by the U.S National Science Foundation under Grants No. DEB 07-20664 and DEB-09-55849. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the NSF. Max Moulds and an anonymous reviewer made suggestions to improve the manuscript.

**References**

Ahmed Z, Sanborn AF (2010) The cicada fauna of Pakistan including the description of four new species (Hemiptera: Cicadoidea: Cicadidae). *Zootaxa* 2516: 27–48.  
 Ahmed Z, Sanborn AF, Hill KBR (2010) A new species of the cicada genus *Cicadatra* from Pakistan (Hemiptera: Cicadoidea: Cicadidae). *Zoology in the Middle East* 51: 75–81.  
 Boulard M (1977) Description d’une nouvelle *Tibicina* de l’ouest Asiatique; révision de la liste des espèces paléartiques appartenant à ce genre (Homoptera: Cicadoidea). *Annales de la Société entomologique de France* 12: 557–66.  
 China WE (1926) A new species of *Cicadatra* (Homoptera, *Cicadidae*) from Waziristan, with notes on the allied genus *Psalmocharias*, Kirk. *Annals and Magazine of Natural History* (9) 18: 374–376. doi: 10.1080/00222932608633531  
 Distant WL (1888) Descriptions of new species of Oriental Cicadidae. *Annals and Magazine of Natural History* (6) 1: 370–376. doi: 10.1080/00222938809460747  
 Distant WL (1904) Additions of a knowledge of the family Cicadidae. *Transaction of the Royal Entomological Society of London* 1904: 667–676.  
 Distant WL (1906) Some undescribed species of Cicadidae. *Annals and Magazine of Natural History* (7) 17: 182–185. doi: 10.1080/00222930608562510

- Dlabola J (1960) Iranische zikaden (Homoptera, Auchenorrhyncha) (Ergebnisse der Entomologischen reisen Willi Richter, Stuttgart, in Iran 1954 und 1956 - nr. 31). Stuttgarter Beitrage zur Naturkunde aus dem Staatlichen Museum für Naturkunde in Stuttgart 41: 1–24.
- Dlabola J (1970) Beitrag zur taxonomie und chorologie einiger palaearktischer zikadenarten (Homoptera, Auchenorrhyncha). Mitteilungen der Münchner Entomologischen Gesellschaft 59: 90–107.
- Dlabola J (1979) *Babuflata* gen.n., neue Membraciden-und Cicadiden-Arten aus dem Iran. Reichenbachia, Staatliches Museum für Tierkunde Dresden 17: 229–241.
- Dlabola J (1981) Ergebnisse der Tschechoslovakisch-Iranische entomologischen expeditionen nach dem Iran 1970 und 1973 (Homoptera: Auchenorrhyncha), II teil. Sborník faunistických prací. Acta Entomologica Musei Nationalis Pragae 40: 127–311.
- Dlabola J, Heller F (1962) Iranische zikaden II. (Ergebnisse der entomologischen Reisen Willi Richter, Stuttgart, im Iran 1954 und 1956 - Nr. 42). Stuttgarter Beitrage zur Naturkunde aus dem Staatlichen Museum für Naturkunde in Stuttgart 90: 1–8.
- Kirkaldy GW (1909) Hemiptera, old and new, No. 2. Canadian Entomologist 41: 388–392. doi: 0.4039/Ent41388-11
- Kolenati F (1857) Homoptera Latreille. Leach. Gulaerostria Zetterstedt. Bulletin de la Société Impériale des Naturalistes Moscuo, Section Biologique 30: 399–444.
- Linnavuori R (1962) Hemiptera of Israel. III. Annales Zoologici Societatis Zoologicae-Botanicæ Fennicæ 'Vanamo' 24: 1–108.
- Melichar L (1896) Cicadinen (Hemiptera-Homoptera) von Mittel-Europa. Verlag von Felix L. Dames, Berlin, 364 pp. doi: 10.5962/bhl.title.8568
- Metcalf ZP (1963) General catalogue of the Homoptera, Fascicle VIII. Cicadoidea. Part 1. Cicadidae. Section II. Gaeninae and Cicadinae. North Carolina State College Contribution 1502: 587–919.
- Mirzayans H (1995) Insects of Iran. The list of Homoptera: Auchenorrhyncha in the insect collection of Plant Pests & Diseases Research Institute. Plant Pests & Diseases Research Institute, Insects Taxonomy Research Department Publication Number 1, 59 pp.
- Moulds MS (2005) An appraisal of the higher classification of cicadas (Hemiptera: Cicadoidea) with special reference to the Australian fauna. Records of the Australian Museum 57: 375–446. doi: 10.3853/j.0067-1975.57.2005.1447
- Mozaffarian F, Sanborn AF (2010) The cicadas of Iran with the description of two new species (Hemiptera: Cicadoidea: Cicadidae). Mitteilungen aus dem Museum für Naturkunde in Berlin – Deutsche Entomologische Zeitschrift 57: 69–84.
- Mozaffarian F, Sanborn AF, Phillips PK (2010) *Cicadatra lorestanica*, a new species of cicada from Iran (Hemiptera: Cicadidae). Journal of Entomological and Aracological Research 42: 27–37.
- Walker F (1850) List of the specimens of homopterous insects in the collection of the British Museum. British Museum Trustees, London, 260 pp.